




Considering Long Term Drought When Prescribed Burning

Robert Lipford—Staff Forester—FM, NC Forest Service, Clayton, NC
Bill Pickens – Conifer Silviculturist, NC Forest Service, Raleigh, NC


Figure 1—Drought as of September 27, 2016



Drought

The summer of 2016 was extremely dry in western North Carolina and, by fall, drought conditions became extreme. By November 1, 2016, fully 2/3 of the NWS Greenville/Spartanburg forecast area was at the D-3 Extreme Drought level and almost 1/4 at the D-4 Exceptional Drought level (Figure 1 & 2). This drought brought one of the most severe fall fire seasons in NCFS history.

Figure 2— Drought as of November 1, 2016



Prescribed Burning—Benefit vs. Risk

The NCFS promotes understory burning for the purpose of hazard reduction, aesthetics, wildlife habitat improvement, and other silvicultural reasons. All burn bosses understand that burning, while beneficial, contains elements of risk to overstory trees. It is his job to minimize the heat from a prescribed burn so as to reduce the risk of damage to valuable overstory trees. The art and science of burning requires that burn bosses consider temperature, relative humidity, wind speed, fuel type, fuel load, fuel availability, season, burning history, and firing technique to control the intensity and duration of heat. Despite responsible planning, every burn boss has unintentionally scorched tree crowns (Figure 3). Many have seen increased overstory mortality.




Figure 3—Crown Scorch

The Effects of Drought on Trees

“Water is essential for almost every plant function. When drought stress occurs, the plant reacts in many ways.” (Blaedow 2013)

“Transpiration is the movement of water from the soil through the roots, stem, branches, leaves, and finally lost to the atmosphere. It is driven by negative pressure, like water being sucked through a straw. The loss of water from the leaves creates a suction that pulls water all the way through the tree from the soil. Water travels through elongated cells in the xylem (sapwood) that resemble straws. In hardwoods they are called vessels, in conifers they are called tracheids. The important thing to note is that this long column of water in vessels is under significant tension. If the water potential of the soil (the water available to be moved into the roots) becomes too low and the leaves are losing too much water, the water column can actually break. . . . The larger the vessels diameter and the longer the water the column, the more likely it is to break if soil moisture is inadequate.” (Blaedow 2013)

“If the water column in the xylem breaks under this tension, an embolism forms. An embolism is an air bubble in the vessel. Once formed, water can no longer flow upward. This is devastating to the plant. If enough embolisms form in enough of the vessels, the plant will not be able to pull enough water out of the soil to survive. The formation of an embolism is known as “cavitation” or “hydraulic failure.” (Blaedow 2013)

Signs of Physiological Drought Stress

- Wilting Foliage
- Wilting Stems
- Leaf Scorch
- Leaf Spotting or Chlorosis
- Early Fall Coloration
- Leaf Senescence
- Bark Cracking
- Stem Bleeding
- Dieback
- Reduced Growth
- Stunting
- Insect Attack
- Disease Spread

- Mortality

Increased Insect and Disease Due to Drought

“Plants under (drought) stress become more suitable as food for herbivorous insects and pathogens resulting in increased incidence and severity of outbreaks” (Blaedow, 2013)

“Drought stressed trees tend to have foliage that is more rich in nitrogen and soluble sugars, callus formation and compartmentalization are weakened, growth is slowed, resin production is reduced, and stressed trees give off many signals that pests can detect such as volatile compounds (e.g. ethanol and α – Pinene), increased foliage reflectivity, increased temperature, and there is even evidence that some insects can detect the sound of cavitation occurring in the xylem. Also, warmer temperatures during the winter allow greater overwintering survival of insects, and perhaps more generations per year allowing larger populations.” (Blaedow 2013)

“However, there are other plant responses that would not seem to be good for insects and diseases. Drought stressed trees have increased amounts of toxins, especially in the foliage. Wood moisture can become so low it is not suitable as a food source. Insect predator and parasite populations can also increase drastically with warmer temperatures. Changes in climate may result in insect emergence at the wrong time for specific feeding, or emergence may be spread out over a longer period of time so that mass attacks are not effective. Drought stressed trees have smaller, stunted tissues that are not as good of a food source for many insects.” (Blaedow 2013)

In sum, foliage feeders, foliage diseases, and pests that require adequate soil moisture, do not benefit from drought conditions in trees, but sap feeders do. Pests like hypoxylon canker, bark beetles, and wood borers, thrive in drought conditions.

Other Stressors In Trees

Drought is not the only stressor for overstory trees. Other known stressors include:

- Overstocked stands are more stressed than properly stocked stands and tend to attract insects.

- Stands in old fields tend to have more *Ips* beetle problems than ones that were established on cut-over land.
- Pine stands that have been raked for straw tend to have more *Ips* beetle problems than non-raked stands. Both loblolly and longleaf are affected by this.
- Stands are significantly more stressed if there has been prior crown damage from wind or ice storms.
- Stands established on eroded land are more stressed.
- Stands subjected to excess nutrient from agricultural field runoff, or subjected to overspray of agricultural chemicals, are more stressed.

Fire as a Stressor

Fire is a stress on overstory trees, but fire can range from a 6” backing fire to a raging head fire. Low intensity fire can be a minor stress, while intense fire can be terminal.


The burn boss has to apply a burning technique and a forester’s knowledge of tree condition. A healthy stand can easily withstand moderate understory fire, but a stressed stand may suffer from flames of much lower intensity. There are no easy “rules of thumb.”

During the summer of 2011, the Ouachita National Forest in Arkansas experienced a severe drought during which the Keetch Byrum Drought Index (Keetch and Byram 1968) was over 700. The High Peak Wildfire burned almost 1500 acres between 29 July and 11 August 2011. The burn site was monitored for damage over the following two years.

“Following the High Peak Wildfire, which ignited from a lightning strike during drought conditions, midstory and overstory densities were significantly reduced between preburn and 1 year postburn. Overstory mortality 1 year postburn was higher than typical background mortality (Clark and others 2008, Klos and others 2009), but mortality from 1 to 2 years postburn was not significant. Overstory density in hardwood and pine-oak forests was reduced to 267 and 356 stems/ha, respectively, moving them closer to historical stand structures. At the same time, pine plantations experienced little mortality and retained an overstory density of 933 stems/ ha.” (Figure 4) (McDaniel 2016)

This seems to show that, as you would expect, the drought and subsequent fire was harder on natural hardwood and mixed pine-hard wood stands than on

Figure 4— Midstory stem density by community type from preburn to 2 years postburn on plots on the High Peak Wildfire in Arkansas, 2011-2013 (McDaniel, 2016)



managed pine plantations.

Caution

Without doubt, understory burning of sufficient intensity to top-kill understory plants has the potential to significantly stress trees in the overstory. The burn boss can partially mitigate this risk by choosing to burn the tract under “cooler” conditions . . . but only by accepting less understory control and less fuel reduction.

Before burning, the burn boss must do a complete size-up and determine if the stand is healthy enough to endure a moderate understory burn. This size-up should consider all stressors and all risks.

Long-term drought adds one more risk factor to these burns.

Two Case Studies from the NCFS Forest Health Branch

Rob Trickle head of the NCFS Forest Health Branch reported on two stands that were burned following a drought that had drastically different results.

“Two years ago, Wayne Langston and I were called by a consultant in Moore County who had recently conducted a late winter/spring prescribed burn that became a hot monster head-fire in a very mature, well managed (BA ≈40-50 sqft/ac) and frequently burned longleaf stand. The consultant was sure he had created beetle bait and neighboring *Ips* beetle populations would pounce on the stressed trees. This stand may have had a few trees attacked by *Ips* beetle over the years, but it was not a problem on this property—however, post drought *Ips* beetle outbreaks were abundant in the area. In the hot spots, understory longleaf pines were killed, some totally consumed. Mature trees were singed to the top. On the 50 or so acres we looked at, buds were breaking on the mature trees and the new candles looked healthy. Immature longleaf were also recovering where the fire was not as hot. We did not find any indication of the presence of *Ips* beetle on the property. Instead, we found 19 trees (mostly red cockaded woodpecker den trees and ones that were previously cat-faced for turpentine production) that were hit hard by black turpentine

beetle. Sap-flow in the larger trees was healthy as indicated by typical pitch tubes. We told the consultant to stand down and observe—we did not think *Ips* beetle would be a problem—so far this assessment has been correct and very few mature trees have been lost.”

“Last year, we were alerted to an *Ips* beetle outbreak in a longleaf stand in Bladen County. The 56-year old stand had been thinned in 2012 (at the end of the drought) and burned in the fall 2015 (flame lengths were re-ported to be very low as the fire backed through the stand). It should be noted that the pre-thinned BA for this stand was 220 sq.ft./ac. and the stand was thinned down to 90.

When Wayne Langston visited the stand, almost half of the standing trees were dead. Some of the green trees were infested with *Ips* beetle and/or ambrosia beetles. Pitch tubes were lacking or very small, an indicator of heavy stress. The fire was blamed, but the situation was nuanced. Another nearby stand that had been thinned at the same time and burned in 2010 and 2014 had similar mortality. Both stands (totaling 54 acres) were salvaged.”

What caused the problem in this second example that resulted in an *Ips* beetle outbreak? Was it stress caused by the initial stocking? Short-term shock from thinning? Burning? Prolonged drought? Age? Likely, the answer is the combination of all of these stresses. This is known as cumulative stress—no one of the factors that lead to mortality held the ‘smoking gun’, but all were contributors.”



Figure 5—Crown Scorch

As Rob Trickel says:

“I would intuitively go lightly (if at all) on the management (thinning and burning) during and directly after a heavy drought for stressed, unmanaged, or poorly managed commercial and overmature stands. I would still be very cautious with management practices that cause short-term stress during, or right after, prolonged drought in well managed stands, but there is more leeway as healthy stands can withstand stress conditions better than stands that are already stressed. It’s hard to make a blanket statement about management during stressful conditions because there are so many factors involved.”

References

- Blaedow, Ryan, A. “How Trees Respond to Stress—The Interactive Effects of Drought and Forest Pests”, PowerPoint Presentation, Written February 10, 2013
- Clark, S.; Spetich, M.; Evans, Z. 2008. Drought in the Southeast. *Forest Wisdom*. 12: 4-13.
- Desperez-Loustau, M. et. al. 2006. Interactive effects of drought and pathogens in forest trees. *Annals of Forest Science* 63: 597-612
- Hsiao, T.C. et. al. 1976. Water stress, growth, and osmotic adjustment. *Phil. Trans. R. Soc. London* 273: 479-500
- Huberty, A.F. and Denno, R.F. 2004. Plant water stress and its consequences for herbivorous insects: a new synthesis. *Ecology* 85: 1383-1398
- Jactel, H. et. al. 2011. Drought effects on damage by forest insects and pathogens: a meta analysis. *Global Climate Change Biology*
- Klos, R.J.; Wang, G.G.; Bauerle, W.L.; Rieck, J.R. 2009. Drought impact on forest growth and mortality in the southeast USA: an analysis using Forest Health and Monitoring data. *Ecological Applications*. 19(3): 699-708.
- Larrson, S. 1989. Stressful times for the plant stress: insect performance hypothesis. *Oikos* 56:277-283
- Mattson, W.J. and Haack, R.A. 1987. The role of drought in outbreaks of plant-eating insects. *Bioscience* 37: 110-118
- McDaniel, Virginia L.; Guldin, James M.; Koerth, Nancy E.; Milks, Jason E.; Finzer, Rebecca J.; Rowland, Ben F. 2016. “Tree Mortality Following a Drought-Year Lightning Ignition in The Ouachita Mountains, Arkansas: 2 Years Postburn.” *Proceedings of The 18th Biennial Southern Silvicultural Research Conference*. Pg. 206-213
- McDowell, N. et. al. 2008. Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New Phytologist* 178: 719-739
- <http://droughtmonitor.unl.edu/MapsAndData/DataTables.aspx>



Figure 6 -Pitch tube from Ips beetle seen in this photo may not form during a drought.

UGA0745096